

Abstract for Third Annual Conference on Carbon Capture and Sequestration
NETL

May 2–6, 2004
Alexandria, VA

Geologic CO₂ Storage Verification Monitoring: Preliminary Results for the Frio Pilot Test

C.M. Oldenburg², and J.L. Lewicki

Earth Sciences Division 90-1116
Lawrence Berkeley National Laboratory
Berkeley CA 94720

Cmoldenburg@lbl.gov; (510) 486-7419; fax: (510) 486-5686
Jllewicki@lbl.gov; (510) 495-2818; fax: (510) 486-5686

²Presenter

We are carrying out baseline measurements of CO₂ concentration and flux in the near-surface environment at South Liberty, Texas, site of a proposed CO₂ injection pilot test. The plan is to inject 3750 tonnes of CO₂ into the Frio Formation at a depth of approximately 1500 m (5000 ft). Pre-injection CO₂ monitoring at the site will provide a baseline against which post-injection CO₂ measurements can be compared to verify that injected CO₂ has not leaked from the deep Frio sands to the shallow subsurface from which it can seep out of the ground. Monitoring of CO₂ fluxes and concentrations is complicated by numerous natural carbon-cycle processes that produce CO₂ flux and concentration variations that can be larger than the CO₂ leakage or seepage signal (CO₂ LOSS). We propose an integrated measurement, monitoring, and modeling strategy that emphasizes cost- and time-efficient measurements designed to discern a CO₂ LOSS from background natural variability. We are using the accumulation chamber (AC) method to measure CO₂ flux at the ground surface, and soil-gas probes to collect gas samples at several depth intervals for CO₂ concentration and carbon isotopic determinations. A later phase of flux measurement is planned using the eddy correlation (EC) approach to capture the effects of the tree canopy on the total CO₂ budget. Preliminary pre-injection results will be presented and discussed.